



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**Parus bicolor.** TUFTED TITMOUSE.—Everywhere common up to about 4000 feet elevation. At Bakersville and Cranberry I seldom took a stroll without hearing the whistle of this bird or seeing it with its young.

**Parus carolinensis.** CAROLINA CHICKADEE.—This bird was not frequently seen, and not observed at all above 5000 feet altitude.

**Regulus satrapa.** GOLDEN-CRESTED KINGLET.—On July 23, shot a young of the year at an elevation of 6000 feet; it was in company with the rest of the brood and the parents. Not common.

**Merula migratoria.** ROBIN.—Rarely seen in the lowlands, but common on the summit of Roan Mountain, where I found two nests with eggs in the balsams; altitude 6300 feet; others were brought to me which had been taken in the woods far from any habitation.

---

## DESCRIPTION OF A NEW *EUETHIA* FROM OLD PROVIDENCE ISLAND.

BY CHARLES B. CORY.

### *Euethia grandior*, sp. nov.

SP. CHAR. —General appearance of *E. bicolor*; but larger, the wing is much longer, and both mandibles are black.

*Adult* ♂ (Type, No. 10,274, Coll. C. B. Cory):—Head, throat, breast, and upper belly black; a patch of dull yellowish white from the belly to the vent; flanks and under tail-coverts dull olive green; back and rump olive green; quills brown, edged with olive green; tail greenish above, slaty brown beneath, and showing numerous indistinct bands when held in the light; bill black.

Length, 4.50; wing, 2.45; tail, 2; tarsus, .60; bill, .40.

The great length of wing seems to be constant in sixteen specimens from Old Providence. I have compared it with forty-four examples of *E. bicolor* from other localities.

*Habitat.* Island of Old Providence, Caribbean Sea.

---

## RECENT LITERATURE.

**The New Canadian Ornithology.\***—Decidedly the best we have is this

---

\*The | Birds of Ontario, | Being a list of Birds observed in the Province of | Ontario, with an Account of their Habits, | Distribution, Nests, Eggs, &c., | —By— | Thomas McIlwraith, | Superintendent of the Ontario District | for the | Migration Committee of the American | Ornithologists' Union | — | Published by the Hamilton Association. | — | Hamilton: | A. Lawson & Co., Printers, 10 York Street. | — | 1886. 1 vol. 8vo. pp. 1-304, i-iv, i-iv, i-vii.